Calcaneal Apophysitis (Sever's Disease)




466

What Is Calcaneal Apophysitis?

Calcaneal apophysitis is a painful inflammation of the heel's growth plate. It typically affects children between the ages of 8 and 14 years old, because the heel bone (calcaneus) is not fully developed until at least age 14. Until then, new bone is forming at the growth plate (physis), a weak area located at the back of the heel. When there is too much repetitive stress on the growth plate, inflammation can develop.

Calcaneal apophysitis is also called Sever's disease, although it is not a true "disease." It is the most common cause of heel pain in children, and can occur in one or both feet.

Heel pain in children differs from the most common type of heel pain experienced by adults. While heel pain in adults usually subsides after a period of walking, pediatric heel pain generally doesn't improve in this manner. In fact, walking typically makes the pain worse.



Causes

Overuse and stress on the heel bone through participation in sports is a major cause of calcaneal apophysitis. The heel's growth plate is sensitive to repeated running and pounding on hard surfaces, resulting in muscle strain and inflamed tissue. For this reason, children and adolescents involved in soccer, track, or basketball are especially vulnerable.

Other potential causes of calcaneal apophysitis include obesity,

a tight Achilles tendon, and biomechanical problems such as flatfoot or a high-arched foot.

Symptoms

Symptoms of calcaneal apophysitis may include:

- Pain in the back or bottom of the heel
- Limping
- Walking on toes
- Difficulty running, jumping, or participating in usual activities or sports
- Pain when the sides of the heel are squeezed

Diagnosis

To diagnose the cause of the child's heel pain and rule out other more serious conditions, the foot and ankle surgeon obtains a thorough medical history and asks questions about recent activities. The surgeon will also examine the child's foot and leg. X-rays are often used to evaluate the condition. Other advanced imaging studies and laboratory tests may also be ordered.

Treatment

The surgeon may select one or more of the following options to treat calcaneal apophysitis:

- **Reduce activity.** The child needs to reduce or stop any activity that causes pain.
- **Support the heel.** Temporary shoe inserts or custom orthotic devices may provide support for the heel.
- Medications. Nonsteroidal anti-inflammatory drugs (NSAIDs), such as ibuprofen, help reduce the pain and inflammation.
- Physical therapy. Stretching or physical therapy modalities are sometimes used to promote healing of the inflamed issue.
- **Immobilization.** In some severe cases of pediatric heel pain, a cast may be used to promote healing while keeping the foot and ankle totally immobile.

PATIENT EDUCATION

Calcaneal Apophysitis continued




Often heel pain in children returns after it has been treated because the heel bone is still growing. Recurrence of heel pain may be a sign of calcaneal apophysitis, or it may indicate a different problem. If your child has a repeat bout of heel pain, be sure to make an appointment with your foot and ankle surgeon.

Can Calcaneal Apophysitis Be Prevented?

The chances of a child developing heel pain can be reduced by:

- Avoiding obesity
- Choosing well-constructed, supportive shoes that are appropriate for the child's activity
- Avoiding or limiting wearing of cleated athletic shoes
- Avoiding activity beyond a child's ability.



This information has been prepared by the Consumer Education Committee of the American College of Foot and Ankle Surgeons, a professional society of over 6,000 foot and ankle surgeons. Members of the College are Doctors of Podiatric Medicine who have received additional training through surgical residency programs.

The mission of the College is to promote superior care of foot and ankle surgical patients through education, research and the promotion of the highest professional standards.